

Diagnostic accuracy of the FINDRISC questionnaire for detecting type 2 diabetes mellitus in an indigenous population from Northeastern Sierra of Puebla, Mexico

Maylin Almonte-Becerril^a , América Martínez-Calleja^b , Nancy Marbella Parra-Torres^a , Geu Mendoza-Catalán^{c*} 

^aDirección ejecutiva de Investigación y de Estudios Avanzados, Universidad de la Salud, Ciudad de México, México; ^bUniversidad para el Bienestar Benito Juárez García, Estado de Guerrero, México; ^cUnidad Académica de Enfermería, Universidad Autónoma del Estado de Hidalgo, Pachuca, México

ABSTRACT

INTRODUCTION The Finnish Diabetes Risk Score questionnaire is a widely used tool for screening type 2 diabetes mellitus; however, its application in Indigenous populations has been little studied. This study assessed the diagnostic accuracy of this questionnaire for detecting cases consistent with type 2 diabetes mellitus in the Totonac population of the state of Puebla, Mexico, and its performance when stratified by gender.

METHODS A cross-sectional diagnostic accuracy study with prospective data collection was conducted in 148 adults (≥ 18 years) without a previous diagnosis of type 2 diabetes mellitus. Participants were residents of communities in the Sierra Nororiental of Puebla and were recruited through convenience sampling with consecutive enrollment during a health fair. The Finnish Diabetes Risk Score questionnaire was used as the index test, evaluated using two cutoff points (≥ 12 and ≥ 15); glycated hemoglobin ($\geq 6.5\%$) was used as the reference standard. Both tests were administered independently and with assessor blinding. Sensitivity, specificity, predictive values, diagnostic accuracy, and the area under the curve were estimated through overall and gender-stratified analyses.

RESULTS At the ≥ 12 cut-off point, the Finnish Diabetes Risk Score showed an overall sensitivity of 76.4% (95%; CI: 66.6 to 84.0), specificity of 55.9% (95%; CI: 43.3 to 67.9), diagnostic accuracy of 68.2% (95%; CI: 60.4 to 75.2), and an area under the curve of 0.64 (95%; CI: 0.54 to 0.73). In women, sensitivity reached 85.2% (95%; CI: 73.4 to 92.3) and the area under the curve was 0.71 (95%; CI: 0.60 to 0.81), whereas in men the discriminative capacity was limited.

CONCLUSIONS The Finnish Diabetes Risk Score questionnaire demonstrated moderate discriminative capacity for detecting cases compatible with type 2 diabetes mellitus in the Totonac Indigenous population, with acceptable performance in women using the cut-off point of ≥ 12 . These findings support its use as an initial screening tool in settings with limited access to confirmatory tests and highlight the need for validation in specific contexts.

KEYWORDS Type 2 diabetes mellitus, indigenous communities, glycated hemoglobin, FINDRISC, diagnostic accuracy, diabetes mellitus tipo 2, hemoglobina glicosilada, comunidades indígenas, desempeño diagnóstico, findrisc

* Corresponding author geu_mendoza@uaeh.edu.mx

Citation Almonte-Becerril M, Martínez-Calleja A, Parra-Torres NM, Mendoza-Catalán G. Diagnostic accuracy of the FINDRISC questionnaire for detecting type 2 diabetes mellitus in an indigenous population from Northeastern Sierra of Puebla, Mexico. Medwave 2026;26(07):e3195
DOI 10.5867/medwave.2026.07.3195

Submitted Dec 17, 2025, **Accepted** Jul 8, 2026,

Published Aug 3, 2026

Postal address Mariano Abasolo 600, Colonia Centro, Pachuca de Soto, Hidalgo 42000, México

INTRODUCTION

Type 2 diabetes mellitus is a chronic disease considered a global public health emergency due to the sustained increase in its prevalence, especially in developing countries [1]. According to the International Diabetes Federation (2025), the global prevalence is 11%, corresponding to 589 million people aged 20 to 79; of these, at least four out of ten remain undiagnosed [2]. In Mexico, the prevalence of type 2 diabetes mellitus and prediabetes is 18.3%, and that of prediabetes is 22.1%. In rural areas, these prevalence rates are 15.2% and 22.8%, respectively [3,4].

MAIN MESSAGES

- Type 2 diabetes mellitus poses a significant burden on indigenous populations, where access to timely diagnosis is limited and there are few validated screening tools available.
- This study evaluated the diagnostic accuracy of the Finnish Diabetes Risk Score questionnaire in a Totonac indigenous population, using glycated hemoglobin as the gold standard, providing the first evidence on its performance in this community.
- Some limitations of this study include potential selection bias due to the voluntary nature of participant enrollment; diagnostic estimates whose accuracy may be compromised in analyses stratified by gender, given that no formal sample size calculation was performed before the study; and a limited evaluation of the instrument's performance in specific subgroups due to the absence of complementary variables.

The magnitude of the burden that diabetes places on health care services and the economic cost it entails for institutions, governments, and families has led to the implementation of specific strategies [5]. In Mexico, the Official Mexican Standard NOM-015-SSA2-2010 for the prevention, treatment, and control of diabetes mellitus remains the current regulatory benchmark [6]. This standard emphasizes the timely detection of the disease and its complications, as well as the implementation of preventive measures among vulnerable groups.

In addition, the current diagnostic criteria for type 2 diabetes mellitus include an oral glucose tolerance test and glycated hemoglobin of 6.5% or higher [7]. However, many public institutions still face limitations related to the availability of supplies, infrastructure, and trained personnel to perform these tests [8,9]. This limitation is even more pronounced in indigenous communities, where additional structural barriers exist, such as limited access to healthcare services, reduced availability of educational resources, and economic constraints [10].

In this context, it is necessary to have accessible, low-cost, and easy-to-implement screening tools that enable the timely identification of cases consistent with diabetes in vulnerable populations [5,11]. Among indigenous populations, two instruments have been primarily used for this purpose: the American Diabetes Association's Diabetes Risk Calculator and the Finnish Diabetes Risk Score (FINDRISC) questionnaire [12–14]. This questionnaire has been widely used internationally [15,16] and has been validated in urban areas of Mexico [13,17]. Although it was originally developed to estimate the future risk of type 2 diabetes mellitus, it has been used as a screening tool to identify undiagnosed cases [16,18]. However, evidence regarding its diagnostic performance in the Mexican indigenous population—particularly in comparison with biochemical tests such as glycated hemoglobin—is limited [12,19,20].

A recent systematic review and meta-analysis reported that the sensitivity and specificity of the Finnish Diabetes Risk Score vary substantially depending on the cutoff point and the population context [18]. This supports the need to evaluate its performance in specific epidemiological and sociocultural contexts [15,18,20]. In this regard, the main objective of the present study was to evaluate the diagnostic accuracy of the Finnish Diabetes Risk Score questionnaire, at cutoff points of 12

or higher and 15 or higher, for the detection of type 2 diabetes mellitus in the Totonac population of the state of Puebla, using glycated hemoglobin as the reference standard, and to assess its performance through a gender-stratified analysis.

METHODS

Design

A cross-sectional diagnostic accuracy study with prospective data collection was conducted, in which the Finnish Diabetes Risk Score questionnaire was used as the index test and glycated hemoglobin as the reference standard. Both tests were administered during the same community outreach event using independent and consecutive procedures, while maintaining blinding between the evaluation teams. The study design, conduct, and reporting were carried out in accordance with the Standards for Reporting of Diagnostic Accuracy (STARD) guidelines [21].

Study sample

The sample consisted of 148 adults aged 18 years or older, both men and women, from the communities of Ixtepec and Huehuetla, Puebla, Mexico. Participants were recruited through convenience sampling with consecutive inclusion during a community health fair organized by the Intercultural University of the State of Puebla. The inclusion criteria were: being 18 years of age or older, agreeing to participate by signing an informed consent form, and having no prior diagnosis of type 2 diabetes mellitus. Participants who did not complete both diagnostic tests were excluded.

Furthermore, due to the operational characteristics of the community health event and the nature of the indigenous population studied, no formal sample size calculation was performed before the study. Therefore, all eligible participants during the data collection period were included.

Diagnostic tools and tests

The Spanish version [22] of the Finnish Diabetes Risk Score questionnaire, based on the original instrument [23], was used as the index test. This test includes eight components: age, body-mass index, waist circumference, physical activity level, fruit and vegetable intake, use of antihypertensive medications,

history of hyperglycemia, and family history of type 2 diabetes mellitus. Each component contributes a specific score that generates the instrument's total score. For this study, cutoff points of 12 or higher and 15 or higher were used; these were defined a priori based on previous reports evaluating the performance of the Finnish Diabetes Risk Score [15,18,20].

A cutoff point of 12 or higher has shown greater sensitivity in population-based screening studies and in Latin American contexts, while a cutoff point of 15 or higher corresponds to the traditional cutoff used for this instrument.

Glycated hemoglobin was used as the reference standard because it reflects the average blood glucose level over the past two to three months without requiring prior fasting and exhibits lower intra-individual biological variability [24,25]. It was measured using a capillary blood sample with the Eclipse A1c™ portable system (Kabla Comercial S.A. de C.V., Mexico). The device was calibrated daily according to the manufacturer's operating instructions before measurements began, and the tests were performed by previously trained personnel. A glycated hemoglobin value of 6.5% or higher was considered indicative of type 2 diabetes mellitus [7,26].

Operating procedure and blinding

The Finnish Diabetes Risk Score questionnaire and the glycated hemoglobin test were administered on the same day through separate workstations, with intervals of less than 30 minutes between each test. Initially, all participants who met the eligibility criteria were assigned a unique, sequential identification code. In addition, they were enrolled consecutively as they arrived at the health fair.

A first team administered the Finnish Diabetes Risk Score questionnaire through a face-to-face interview in Spanish. For participants with limited proficiency in Spanish, bilingual staff facilitated understanding of the items by providing verbal explanations in their native language (Tutunakú), without altering the original structure or content of the instrument. Responses were recorded in Spanish for analysis.

Subsequently, participants were directed to a second station, where glycated hemoglobin samples were collected and processed using the same coding system. The evaluators did not have access to the results obtained during the fieldwork, ensuring blinding between the evaluation teams. No adverse effects resulting from the administration of the tests were reported.

Data processing and analysis

Statistical analyses were performed using SPSS v.25 (IBM Corp., Armonk, NY, USA). Categorical variables were described using frequencies and percentages, and quantitative variables as mean $\pm$ standard deviation.

Scores obtained from the Finnish Diabetes Risk Score questionnaire at both cutoff points were classified as positive or negative according to each threshold. Subsequently, 2x2 contingency tables were constructed between the results

of the index test and the reference standard. Based on these, sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) were calculated. Likewise, diagnostic accuracy, defined as the total proportion of correct classifications relative to the reference standard, was calculated using the formula $(TP + TN) / \text{Total } N$, where TP corresponds to true positives and TN to true negatives. The 95% confidence intervals were estimated using Wilson's method for binomial proportions [27].

The overall discriminatory ability of the instrument used was evaluated using receiver operating characteristic curves, comparing its continuous score against glycated hemoglobin (6.5% or higher versus less than 6.5%); the area under the curve, estimated using nonparametric methods, served as the measure of overall discrimination.


To characterize the distribution of the disease, participants with glycated hemoglobin of 6.5% or higher were classified according to the severity of their hyperglycemia (6.5 to 6.9%; 7.0 to 7.9%; 8.0 to 8.9%; 9.0% or higher). Meanwhile, participants with glycated hemoglobin below 6.5% were characterized based on relevant metabolic conditions, including nutritional status and the presence of abdominal adiposity. Nutritional status was classified using body-mass index into normal weight (18.5 to 24.9 kilograms per square meter), overweight (25.0 to 29.9 kilograms per square meter), and obese (30.0 kilograms per square meter or higher). Abdominal adiposity was determined by waist circumference and was considered present when it was 80 centimeters or greater in women and 90 centimeters or greater in men [28].

Finally, given that various authors have documented differences in the distribution of risk factors and health behaviors between men and women [18,20,29], a gender-stratified analysis of these same parameters was conducted. No additional stratified analyses were performed, as they were not part of the study's objectives and the available sample size did not allow for estimates with sufficient precision in other subgroups. No missing data were identified in the analyzed variables.

RESULTS

During the recruitment period, 245 potentially eligible participants were identified. Of these, 97 were excluded (34 because of having a diagnosis of type 2 diabetes mellitus and 63 because they had not completed both diagnostic tests; Figure 1).

The study sample consisted of 148 adults, predominantly women (66.2%; 98/148) and speakers of Tutunakú (81.7%; 121/148). The average age was 54.3 ± 16.7 years, with a significant difference of 10.6 years between men and women (95% confidence interval: 5.1 to 16.1). Regarding educational attainment, 38.5% (57/148) had completed primary school, while 35.1% (52/148) had no formal education, with a higher percentage among women (38.8%; 38/98). Furthermore, participants had a body-mass index of 26.4 (standard deviation

Figure 1. Flowchart of participant selection, administration of the index test (FINDRISC), and the reference standard (glycated hemoglobin).

FINDRISC: Finnish Diabetes Risk Score; STRARD: Standards for Reporting of Diagnostic Accuracy.

Source: Prepared by the authors based on the study results in accordance with the STARD guidelines [21].

± 3.9) kilograms per square meter and an average waist circumference of 92.6 centimeters (standard deviation ± 8.5) (Table 1).

After describing the sociodemographic characteristics, the participants were classified according to the categories of the index test and the reference standard. In the Finnish Diabetes Risk Score classification, 44.9% (44/98) of the women and 22.0% (11/50) of the men were classified as high risk, while 33.7% (33/98) of the women and 42.0% (21/50) of the men were classified as low risk. Regarding glycated hemoglobin levels, only 5.4% (8/148) had values consistent with normoglycemia, and 34.5% (51/148) had levels consistent with prediabetes, with the latter being higher among women. In contrast, 55.1% (54/98) of women and 70.0% (35/50) of men had values consistent with type 2 diabetes mellitus (Table 2).

Severity of the condition and relevant metabolic conditions

Subsequently, participants with glycated hemoglobin levels consistent with type 2 diabetes mellitus (60.1%; 89/148) were classified according to glycemic severity. In the total sample, 41.6% (37/89) had glycated hemoglobin values between 7.0% and 7.9%, while 24.7% (22/89) had levels of 8.0% or higher. When analyzing the distribution by gender, a higher proportion of men was observed in the 7.0 to 7.9% category (48.6%; 17/35) compared with women (37.0%; 20/54). In contrast, women had a slightly higher proportion of values equal to or greater than 8.0% (27.8%; 15/54) compared to men (20.0%; 7/35) (Table 3).

In addition, the metabolic conditions present in participants without type 2 diabetes mellitus (glycated hemoglobin < 6.5%) were described. The results showed that prediabetes was the predominant metabolic condition (86.4%; 51/59), of whom 45.1% (23/51) were of normal weight and 43.1%

Table 1. Sociodemographic and clinical characteristics of the study sample.

Variables	Total (N = 148) n (%)	Female (N = 98) n (%)	Male (N = 50) n (%)
Language			
Spanish	27 (18.3)	17 (17.3)	10 (20.0)
Tutunakú	121 (81.7)	81 (82.7)	40 (80.0)
Marital status			
Single	17 (11.5)	9 (9.2)	8 (16.0)
In a relationship	131 (88.5)	89 (90.8)	42 (84.0)
Education level			
No formal education	52 (35.1)	38 (38.8)	14 (28.0)
Elementary school	57 (38.5)	34 (34.7)	23 (46.0)
High school	20 (13.5)	14 (14.3)	6 (12.0)
Vocational school	19 (12.8)	12 (12.2)	7 (14.0)
Clinical characteristics			
	Mean ± SD	Mean ± SD	Mean ± SD
Age (years)	54.3 ± 16.7	50.7 ± 16.9	61.3 ± 14.0
BMI (kg/m ²)	26.4 ± 3.9	26.7 ± 4.0	25.8 ± 3.6
Waist circumference (cm)	92.6 ± 8.5	92.5 ± 8.7	92.7 ± 8.2

BMI, body-mass index. SD, standard deviation. kg, kilograms. m², square meters. cm, centimeters.

Notes: The table includes a breakdown by gender (n = 148).

Source: Prepared by the authors based on the study results.

Table 2. Classification according to FINDRISC and HbA1c in the total sample and by gender.

Classification categories	Total (n = 148) n (%)	Female (n = 98) n (%)	Male (n = 50) n (%)
FINDRISC test			
Low	54 (36.5)	33 (33.7)	21 (42.0)
Moderate	39 (26.4)	21 (21.4)	18 (36.0)
High	55 (37.2)	44 (44.9)	11 (22.0)
Glycemic classification based on HbA1c			
Normal glucose levels (< 5.7%)	8 (5.4)	5 (5.1)	3 (6.0)
Prediabetes (5.7 to 6.4)	51 (34.5)	39 (39.8)	12 (24.0)
Consistent with T2D (≥ 6.5%)	89 (60.1)	54 (55.1)	35 (70.0)

DT2, type 2 diabetes mellitus; HbA1c, glycated hemoglobin; FINDRISC, Finnish Diabetes Risk Score.

Source: Prepared by the authors based on the study results.

Table 3. Distribution of glycemic severity in participants with type 2 diabetes and relevant metabolic conditions in participants without type 2 diabetes.

Distribution of glycemic severity among participants with T2D (N = 89)			
HbA1c level	Total (n = 89) n (%)	Females (n = 54) n (%)	Males (n = 35) n (%)
6.5 to 6.9%	30 (33.7)	19 (35.2)	11 (31.4)
7.0 to 7.9%	37 (41.6)	20 (37.0)	17 (48.6)
8.0 to 8.9%	10 (11.2)	7 (13.0)	3 (8.6)
≥ 9.0%	12 (13.5)	8 (14.8)	4 (11.4)
Relevant metabolic conditions in participants without T2D (HbA1c < 6.5%).			
Variables	Total (n = 59) n (%)	Normal blood glucose (n = 8) n (%)	Prediabetes (n = 51) n (%)
Nutritional status			
Normal weight	25 (42.4)	2 (25.0)	23 (45.1)
Overweight	26 (44.1)	4 (50.0)	22 (43.1)
Obesity	8 (13.5)	2 (25.0)	6 (11.8)
Abdominal adiposity¹			
Absent	11 (18.6)	2 (25.0)	9 (17.7)
Present	48 (81.4)	6 (75.0)	42 (82.3)

T2D, type 2 diabetes mellitus; HbA1c, glycated hemoglobin; cm, centimeters.

Notes: ¹Abdominal adiposity: waist circumference ≥ 80 cm in women and ≥ 90 cm in men.

Notes: Participants with T2D: HbA1c ≥ 6.5%; n = 89. Participants without T2D: HbA1c < 6.5%; n = 59.

Source: Prepared by the authors based on the study results.

(22/51) were overweight. Among participants with normal blood glucose levels, 50% (4/8) were overweight. Furthermore,

Table 4. Distribution of FINDRISC questionnaire results compared to the reference standard.

Sample ¹	FINDRISC ² result	Positive HbA1c n (%)	Negative HbA1c n (%)
Total (n = 148)			
FINDRISC ≥ 12	Positive	68 (72.3)	26 (27.7)
	Negative	21 (38.9)	33 (61.1)
FINDRISC ≥ 15	Positive	39 (70.9)	16 (29.1)
	Negative	50 (53.8)	43 (46.2)
Female (n = 98)			
FINDRISC ≥ 12	Positive	46 (70.7)	19 (29.3)
	Negative	8 (24.2)	25 (75.8)
FINDRISC ≥ 15	Positive	32 (72.7)	12 (27.3)
	Negative	22 (40.7)	32 (59.3)
Male (n = 50)			
FINDRISC ≥ 12	Positive	22 (75.9)	7 (24.1)
	Negative	13 (61.9)	8 (38.1)
FINDRISC ≥ 15	Positive	7 (63.6)	4 (36.4)
	Negative	28 (71.8)	11 (28.2)

FINDRISC: Finnish Diabetes Risk Score. HbA1c: glycated hemoglobin.

Notes: ¹The percentages were calculated by row, based on the total for each FINDRISC category and within each subgroup (total sample, women, and men). ²The cells represent true positives, false positives, false negatives, and true negatives.

Notes: Reference standard: HbA1c $\geq 6.5\%$. Data applied to the total sample and stratified by gender, according to the cutoffs of ≥ 12 and ≥ 15 .

Source: Prepared by the authors based on the study results.

when abdominal adiposity was assessed, 81.4% (48/59) of the participants exhibited it, with a higher prevalence among those with prediabetes (82.3%; 42/51) (Table 3).

Sensitivity and specificity of the Finnish Diabetes Risk Score test

Using the cut-off points of the Finnish Diabetes Risk Score questionnaire as a reference, 2x2 contingency tables were constructed against the reference standard (glycated hemoglobin $\geq 6.5\%$), both for the total sample and for the sample stratified by gender. The results showed that, in the total sample, more than 70% of cases with a positive score also had an HbA1c level of 6.5% or higher (true positives), regardless of the cutoff value used. At a cutoff value of 12 or higher, 61.1% (33/54) of participants with a negative questionnaire score also had an HbA1c level below 6.5% (true negatives); this proportion decreased to 46.2% (43/93) at a cutoff of 15 or higher (Table 4).

When stratified by gender, women showed a higher proportion of true positives, particularly at the cutoff of 15 or higher (72.7%; 32/44). Likewise, the cutoff of 12 or higher showed a higher proportion of true negatives (75.8%; 25/33). In contrast, men showed a higher frequency of false negatives, especially at a cutoff of 15 or higher (71.8%; 28/39, Table 4).

Subsequently, when evaluating the diagnostic performance parameters, the results showed that, in the total sample, a cutoff point of 12 or higher achieved a sensitivity of 76.4% (95% confidence interval: 66.6 to 84.0) and a specificity of 55.9% (95% confidence interval: 43.3 to 67.9). In contrast, for a cutoff point of 15 or higher, lower sensitivity was observed (43.8%; 95% confidence interval: 34.0 to 54.1). Diagnostic accuracy was better at a cutoff point of 12 or higher (68.2%; confidence interval: 95%; 60.4 to 75.2; Table 5).

When stratified by gender, women showed higher sensitivity at both cut-off points, particularly at the cut-off of 12 or higher (85.2%; 95% confidence interval: 73.4 to 92.3) compared with men. In contrast, men showed lower sensitivity, especially for the cutoff of 15 or higher (20.0%; 95% confidence interval: 10.0 to 35.9), achieving a specificity of 73.3% (95% confidence interval: 48.1 to 89.1). The highest percentage of diagnostic accuracy was obtained in samples from women at a cutoff point of 12 or higher (72.5%; 95% confidence interval: 63.0 to 80.3) (Table 5).

Analysis using receiver operating characteristic curves showed that the Finnish Diabetes Risk Score questionnaire had moderate discriminatory power in the total sample, with an area under the curve of 0.64 (95% confidence interval: 0.54 to 0.73). When the analysis was stratified by gender, an area under the curve of 0.71 (95% confidence interval: 0.60 to 0.81) was obtained for women and 0.51 (95% confidence interval: 0.31 to 0.69) for men (Figure 2).

DISCUSSION


Diabetes mellitus is one of the leading public health challenges of the twenty-first century [4,30]. Its burden is disproportionately higher among Indigenous populations, where structural inequalities and barriers to healthcare access substantially contribute to diabetes-related complications and premature mortality [8–10,30,31]. Although several screening tools are available, including the Finnish Diabetes Risk Score (FINDRISC), which has been extensively evaluated in different populations worldwide [18,20,32], its performance in Indigenous populations has been insufficiently investigated. Therefore, the present study provides the first evidence regarding its diagnostic performance in a Totonac community.

Table 5. Diagnostic performance parameters of the FINDRISC questionnaire.

Sample	FINDRISC cut-off point	Sensitivity % (95% CI)	Specificity % (95% CI)	PPV % (95% CI)	NPV % (95% CI)	Diagnostic accuracy % (95% CI)
Total (n = 148)	≥12	76.4% (66.6 to 84.0)	55.9% (43.3 to 67.9)	72.3% (62.6 to 80.4)	61.1% (47.8 to 73.0)	68.2% (60.4 to 75.2)
	≥15	43.8% (34.0 to 54.1)	72.9% (60.4 to 82.6)	70.9% (57.9 to 81.2)	46.2% (36.5 to 56.3)	55.4% (47.4 to 63.2)
Females (n = 98)	≥12	85.2% (73.4 to 92.3)	56.8% (42.2 to 70.3)	70.8% (59.0 to 80.4)	75.8% (59.0 to 87.2)	72.5% (63.0 to 80.3)
	≥15	59.3% (46.0 to 71.3)	72.7% (58.2 to 84.0)	72.7% (58.2 to 83.7)	59.3% (46.0 to 71.3)	65.3% (55.5 to 74.0)
Males (n = 50)	≥12	62.9% (46.3 to 76.8)	53.3% (30.1 to 75.2)	75.9% (57.9 to 87.8)	38.1% (20.8 to 59.1)	60.0% (46.2 to 72.4)
	≥15	20.0% (10.0 to 35.9)	73.3% (48.1 to 89.1)	63.6% (35.4 to 84.8)	28.2% (16.5 to 43.8)	36.0% (24.1 to 49.9)

FINDRISC: Finnish Diabetes Risk Score. CI: confidence interval. PPV: positive predictive value. NPV: negative predictive value.
Notes: Data for the cutoffs of ≥ 12 and ≥ 15 in the total sample and stratified by gender.

Figure 2. ROC curves of the FINDRISC questionnaire for identifying elevated HbA1c (≥ 6.5%).



The ROC curves are shown for the total population (A), women (B), and men (C). The area under the curve values are presented with their 95% confidence intervals. The diagonal line represents the performance expected by chance.
AUC: area under the curve; CI: confidence interval; ROC: receiver operating characteristic curve; FINDRISC: Finnish Diabetes Risk Score; HbA1c: glycated hemoglobin.
Source: Prepared by the authors based on the study results.

Our findings showed that the diagnostic performance of the FINDRISC questionnaire improved when a cutoff value of ≥ 12 was applied, achieving moderate discriminatory ability and a sensitivity of **76.4%** (95% confidence interval: 66.6 to 84.0). These findings are consistent with previous studies demonstrating that demographic, clinical, and epidemiological characteristics of the target population influence the performance of the instrument [16,33–35]. Accordingly, adapting the cutoff value may reduce the number of individuals requiring confirmatory testing while improving the identification of previously undiagnosed cases of type 2 diabetes mellitus [15–18].

Similarly, the gender-stratified analysis revealed that the performance of the instrument was not uniform across subgroups. Among women, the ≥ 12 cutoff yielded the highest sensitivity, even exceeding that reported for other screening

tools developed for non-Indigenous Mexican women [36,37]. In contrast, the instrument demonstrated limited performance among men.

These differences may be explained by both biological and sociocultural factors. From a biological perspective, gender-related differences in body composition and adiposity may influence the ability of the anthropometric components of the questionnaire to accurately reflect the underlying metabolic condition. In women, the predominance of peripheral fat distribution may enhance the relationship between adiposity and body-mass index [38,39]. While in men, this relationship may be less consistent because of greater lean body-mass and a higher accumulation of central and visceral adiposity [38–41].

Furthermore, the high prevalence of overweight, obesity, and abdominal adiposity observed among participants without type

2 diabetes mellitus may also have affected the specificity of the instrument. This finding is consistent with the spectrum effect described for screening and risk prediction tests, whereby a high prevalence of anthropometric risk factors reduces the ability of the instrument to discriminate between individuals with and without diabetes, thereby increasing the number of false-positive results [42].

In addition, previous studies have shown that gender norms and stereotypes influence not only how individuals report their medical history and risk factors [29,33,43], but also their perception of disease risk, self-care practices, and healthcare-seeking behaviors [12,33,34]. Consequently, some of the variables included in the FINDRISC questionnaire may not adequately capture the biological and sociocultural determinants of diabetes risk in this population, thereby limiting its diagnostic performance.

Although the FINDRISC questionnaire showed limited performance among men, its diagnostic performance in women represents a particularly relevant finding. This result is especially important considering that women with type 2 diabetes mellitus experience greater cardiovascular risk and poorer clinical outcomes than men [29]. Moreover, among Mexican Indigenous populations, women bear a higher burden of diabetes-related risk factors, a situation further exacerbated by economic dependence, gender inequalities, and barriers to self-care and timely access to healthcare services [8,31,44].

In this context, the FINDRISC questionnaire, using a cutoff value of ≥ 12 , may represent a useful initial screening tool for Indigenous women. Nevertheless, its implementation should be supported by local validation and contextual adaptation, given that perceptions of health and disease in Indigenous communities are strongly influenced by cultural beliefs, worldviews, and community practices [11,30,31,45].

Study limitations

Several limitations should be considered when interpreting the findings of this study. First, participants were recruited on a voluntary basis, which may have introduced selection bias, as individuals who chose to participate may have been more concerned about their health status or already aware of potential diabetes risk factors.

Second, no formal sample size calculation was performed before the study because all eligible participants available during the data collection period were included from Indigenous communities with relatively small populations. This may have affected the precision of the diagnostic accuracy estimates, particularly in the gender-stratified analyses, as reflected by the relatively wide confidence intervals. Therefore, these findings should be interpreted with caution and considered preliminary evidence for this population.

In addition, the absence of complementary variables limited the possibility of evaluating the performance of the instrument in specific subgroups, which could have provided a more detailed characterization of diabetes risk.

Regarding the linguistic component, the FINDRISC questionnaire did not undergo a formal translation and cross-cultural validation process into the Tutunakú language. Although linguistic mediation was provided by bilingual personnel native to the region, facilitating participants' understanding of the questionnaire items, the possibility of differences in interpretation cannot be completely excluded and therefore represents a methodological limitation. Moreover, because some questionnaire items rely on participants' prior knowledge of their health status, limited access to healthcare services may have reduced this knowledge, potentially affecting the accuracy of some responses regardless of the language used.

CONCLUSIONS

The Finnish Diabetes Risk Score demonstrated moderate diagnostic performance in the Totonac Indigenous population, with better performance among women when a cutoff value of ≥ 12 was used, characterized by high sensitivity but low specificity. These findings suggest that the instrument may be useful as an initial screening tool in this population, particularly in settings where access to biochemical diagnostic testing is limited, provided that positive results are confirmed with diagnostic testing.

In contrast, its limited performance among men highlights the need to develop or adapt more sensitive screening tools for this subgroup. Overall, these findings underscore the importance of considering both gender-related differences and the sociocultural context when evaluating and implementing diabetes screening instruments in Indigenous populations.

Contributor roles MAB: Conceptualization, methodology, supervision, formal analysis, writing, review and editing, and original draft preparation. AMC: Formal analysis, writing, and original draft preparation. NMPT: Methodology, investigation, formal analysis, and original draft preparation. GMC: Conceptualization, methodology, formal analysis, and original draft preparation.

Acknowledgments The authors sincerely thank the participating community members for their willingness, trust, and collaboration. We also extend our special appreciation to the nursing students of the Intercultural University of the State of Puebla for their invaluable assistance during data collection, as well as for their commitment and cultural sensitivity throughout the fieldwork. Their support in facilitating communication in the Tutunakú language enabled a respectful and culturally appropriate interaction with the community.

Conflictos de intereses The authors declare that they have no conflicts of interest related to this work.

Funding This work is based on results obtained from a project previously funded by the Program for Teacher Professional Development (PRODEP). The funding covered the initial phase of the project and was not specifically intended for the preparation or publication of this manuscript. Project number: UIEP-CA-07.

Statement on the use of artificial intelligence models ChatGPT was used as a support tool to improve the writing, clarity, and structure of the manuscript. It was not used to generate original scientific content, perform data analysis, or formulate the study conclusions. All content was critically reviewed, validated, and approved by the authors.

Protocol registry This study is neither a clinical trial nor a systematic review; therefore, it was not registered in protocol registries such as ClinicalTrials.gov or PROSPERO.

Ethics The study was approved by the Intercultural University of the State of Puebla (Approval No. UIEP/CE/SE/01/2022). Participation was voluntary, and the confidentiality and anonymity of all collected information were ensured. The study was conducted in accordance with the ethical principles of the Declaration of Helsinki for research involving human participants.

Data sharing statement The data supporting the findings of this study are not publicly available because of confidentiality considerations. However, they are available from the corresponding author upon reasonable request.

Language of submission Spanish.

Peer review and provenance Not commissioned. Externally peer-reviewed by two reviewers and reviewed by the journal's statistical editor. All reviews were done in a double-anonymous modality.

REFERENCES

- Vergauwen J, Monnet F, Van Olmen J, Wouters E, Danhieux K, Desmet L, et al. Evaluating the effectiveness of lifestyle education for individuals at increased risk of type 2 diabetes and cardiovascular diseases (Halt2Diabetes): protocol for a repeated measures study. *BMC Public Health*. 2025;25. <https://doi.org/10.1186/s12889-025-24462-w>
- International Diabetes Federation, IDF. Datos y cifras. International Diabetes Federation, IDF. 2025;Available. <https://idf.org/es/about-diabetes/diabetes-facts-figures/>
- Basto-Abreu A, López-Olmedo N, Rojas-Martínez R, Aguilar-Salinas CA, Moreno-Banda GL, Carnalla M, et al. Prevalencia de prediabetes y diabetes en México: Ensanut 2022. *Salud Publica Mex*. 2023;65: s163–s168. <https://doi.org/10.21149/14832>
- Reyes-García A, Basto-Abreu A, Reyes-Sánchez F, Stern D, Romero-Martínez M, Campos-Nonato I, et al. Prevalencia de diabetes y control glucémico en México, Ensanut 2021-2024. *Salud Publica Mex*. 2025;67: 622–632. <https://doi.org/10.21149/17286>
- Zehra A, Gerstle D, Ali FM, Ali M, Mejia-Lancheros C, Fazli GS. A Scoping Review on Community-based Diabetes Screening Interventions: Paving the Pathway to Early Care and Prevention of Diabetes. *Curr Diab Rep*. 2025;25. <https://doi.org/10.1007/s11892-025-01605-2>
- Secretaría de Salud. NOM-015-SSA2-2010. Para la prevención, tratamiento y control de la diabetes mellitus. Diario Oficial de la Federación [Internet. 2010. https://diariooficial.gob.mx/nota_detalle.php?codigo=5168074&fecha=23/11/2010#gsc.tab=0
- ElSayed NA, McCoy RG, Aleppo G, Balapattabi K, Beverly EA, Briggs Early K, et al. 2. Diagnosis and Classification of Diabetes: Standards of Care in Diabetes—2025. *Diabetes Care*. 2025;48: S27–S49. <https://doi.org/10.2337/dc25-S002>
- In: Consejo Nacional de Evaluación de la Política de Desarrollo Social, CONEVAL. Pobreza y Población Indígena en México [Internet]. Ciudad de México: CONEVAL; 2025. https://www.coneval.org.mx/InformesPublicaciones/InformesPublicaciones/Documents/Pobreza_Poblacion_Indigena.pdf
- Castro-Porras LV, Rojas-Martínez R, Romero-Martínez M, Aguilar-Salinas CA, Escamilla-Núñez C. The Trend in the Prevalence of Diabetes Mellitus in the Mexican Indigenous Population From 2000 to 2018. *AJPM Focus*. 2023;2. <https://doi.org/10.1016/j.focus.2023.100087>
- Claussen C, Papadimos E, Magliano DJ, Hotu C, Monteith H, Shah B, et al. Prevalence of type 2 diabetes among global Indigenous adult populations: a systematic review. *Diabetologia*. 2026;69: 582–599. <https://doi.org/10.1007/s00125-025-06624-y>
- Arellano Sosa AT, Zurita Paucar PA, Sinchi Vivas NM, Barzola Troya BB, Jaramillo Alaleo AE, Alban Rivera SY. Comorbilidad, Adherencia y Calidad de Vida en Personas con Diabetes Tipo 2: Estudio Comparativo en Población Urbana y Rural. *Revista Científica y Académica*. 2025;5: 1262–1283. <https://estudiosyperspectivas.org/index.php/EstudiosyPerspectivas/issue/view/11> <https://doi.org/10.61384/r.c.a.v5i2.1205>
- Mendoza-Catalán G, Parra-Torres NM, Almonte-Becerril M. Risk of chronic diseases in indigenous Totonacs from Mexico short running title: Risk of chronic diseases in indigenous Totonacs. *Appl Nurs Res*. 2022;63: 151543. <https://doi.org/10.1016/j.apnr.2021.151543>
- Morales Ramirez D. Tamizaje para identificar el riesgo a desarrollar Diabetes Mellitus Tipo 2 en adultos mayores de 20 años. *ARCHIVOS DE MEDICINA*. 2023;2: 2–27. <https://doi.org/10.29059/amsem.v2i2.73>
- Lundholm MD. Updated Review of the American Diabetes Association (ADA) Risk Calculator. *Evidence to Action*. 2026;012026. <https://doi.org/10.65357/001c.159088>
- Aldás-Avila A de los Á, Romo-López ÁG. Sensibilidad y especificidad del cuestionario Findrisc como predictor de riesgo para desarrollar diabetes mellitus tipo 2. Una revisión bibliográfica de los últimos 5 años. *MQRInvestigar*. 2024;8: 84–98. <https://doi.org/10.56048/MQR20225.8.1.2024.84-98>
- Yovera-Aldana M, Mezones-Holguín E, Agüero-Zamora R, Damas-Casani L, Uriol-Llanos B, Espinoza-Morales F, et al. External validation of Finnish diabetes risk score (FINDRISC) and Latin American FINDRISC for screening of undiagnosed dysglycemia: Analysis in a Peruvian hospital health care workers sample. *PLoS ONE*. 2024;19: e0299674. <https://doi.org/10.1371/journal.pone.0299674>
- Varela-Vega Y, Roy-García IA, Pérez-Rodríguez M, Velázquez-López L. Certeza diagnóstica del instrumento FINDRISC para identificar resistencia a la insulina en adultos. *Revista Médica del Instituto Mexicano del Seguro Social*. 2023;61: 33–41.
- Acosta-Reyes J, Rodríguez Garrido DP, Acosta Vergara T, Aschner P, Fraga CA, Vazquez-Fernandez A, et al. Finnish diabetes risk score (findrisc) for type 2 diabetes screening compared with the oral glucose tolerance test: A systematic review and meta-analysis of diagnostic test accuracy.

- Diabetes Res Clin Pract. 2025;229: 112480. <https://doi.org/10.1016/j.diabres.2025.112480>
19. Brito Nuñez JD, Cedeño Rondón J, Pérez Arciniega E, Brito Núñez N. Riesgo de diabetes mellitus según el cuestionario Finnish Diabetes Risk Score (FINDRISC) en Indígenas Warao de Barrancas del Orinoco, Monagas, Venezuela. *Rev salud pública*. 2022;24: 1–5. <https://doi.org/10.15446/rsap.v24n4.75956>
 20. Nieto-Martínez R, Barengo NC, Restrepo M, Grinspan A, Assefi A, Mechanick JL. Large scale application of the Finnish diabetes risk score in Latin American and Caribbean populations: a descriptive study. *Front Endocrinol*. 2023;14: 1188784. <https://doi.org/10.3389/fendo.2023.1188784>
 21. Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig L, et al. STARD 2015: an updated list of essential items for reporting diagnostic accuracy studies. *BMJ*. 2015;351. <https://doi.org/10.1136/bmj.h5527>
 22. Soriguer F, Valdés S, Tapia MJ, Esteva I, Ruiz de Adana MS, Almaraz MC, et al. Validación del FINDRISC (FINnish Diabetes Risk Score) para la predicción del riesgo de diabetes tipo 2 en una población del sur de España. *Estudio Pizarra. Medicina Clínica*. 2012;138: 371–376. <https://doi.org/10.1016/j.medcli.2011.05.025>
 23. Lindström J, Tuomilehto J. The Diabetes Risk Score. *Diabetes Care*. 2003;26: 725–731. <https://doi.org/10.2337/diacare.26.3.725>
 24. Kasujja FX, Daivadanam M, Mayega RW, Nuwaha F, Kusolo R, Ekirapa E. Glycated haemoglobin versus fasting plasma glucose for type 2 diabetes point of care screening: a decision model cost-effectiveness analysis. *BMC Health Serv Res*. 2025;25. <https://doi.org/10.1186/s12913-025-12840-4>
 25. National Institute of Diabetes and Digestive and Kidney Diseases, NIDDK. La prueba de A1C y la diabetes - NIDDK. National Institute of Diabetes and Digestive and Kidney Diseases. <https://www.niddk.nih.gov/health-information/informacion-de-la-salud/pruebas-diagnosticas/prueba-a1c-diabetes>
 26. American Diabetes Association ADA. In: ¿Qué es A1C? | American Diabetes Association [Internet]. <https://diabetes.org/es/sobre-la-diabetes/a1c>
 27. Dean A, Sullivan K, Soe M. Diagnostic Test - OpenEpi module for performance evaluation of a diagnostic test. OpenEpi: Open Source Epidemiologic Statistics for Public Health. https://www.openepi.com/Menu/OE_Menu.htm
 28. Chávez-Manzanera EA, Vera-Zertuche JM, Kaufer-Horwitz M, Vázquez-Velázquez V, Flores-Lázaro JR, Mireles-Zavala L, et al. Guía mexicana de práctica clínica para el manejo del sobrepeso y la obesidad en el adulto. *RME*. 2025;12: 16026. <https://doi.org/10.24875/RME.M24000042>
 29. Kautzky-Willer A, Leutner M, Harreiter J. Sex differences in type 2 diabetes. *Diabetologia*. 2023;66: 986–1002. <https://doi.org/10.1007/s00125-023-05891-x>
 30. Morales A, Granda F, Rochina M, Rosas F, Olaya MC, Lomas J. Diabetes mellitus tipo 2 en pueblos indígenas. *INSPILIP*. 2022; 38–44. <https://doi.org/10.31790/inspilip.v6iEspecial.324>
 31. Kovanur Sampath K, Ann-Rong Y, Brownie S. Culturally appropriate care for indigenous people with type 2 diabetes mellitus (T2DM)- a scoping review. *Prim Care Diabetes*. 2025;19: 238–245. <https://doi.org/10.1016/j.pcd.2025.02.008>
 32. Ríos Rodríguez SJ, Gutiérrez Cueva R, Gutiérrez Ayala GIL, Robles Romero MÁ, Gutiérrez Cueva JF. Riesgo de desarrollar diabetes mellitus tipo 2 según escala Finnish Diabetes Risk Score en atención primaria. *Revista Cubana de Medicina General Integral*. 2023;Available. http://scielo.sld.cu/scielo.php?script=sci_abstract&pid=S0864-21252023000200011&lng=es&nrm=iso&tlng=es
 33. Mendoza-Catalán G, Almonte-Becerril M, Jiménez-Orsorio AS, Ángel-García J, Estrada-Luna D, Rodríguez-Santamaría Y, et al. Machismo and Marianismo Associated With the Risk of Type 2 Diabetes in Indigenous Mexican Adults. *J Prim Care Community Health*. 2025;16. <https://doi.org/10.1177/21501319251357500>
 34. Mendoza-Catalán G, Jimenez-Orsorio AS, Angel-García J, Estrada-Luna D, Librado-Gonzalez N, Carvalho-Gomes I. Masculinity and Self-Care Activities Among Mexican Adult Men With Type 2 Diabetes. *SAGE Open Nurs*. 2026;12. <https://doi.org/10.1177/23779608261430134>
 35. de Keijzer B, Cuellar AC, Valenzuela Mayorga A, Hommes C, Caffé S, Mendoza F, et al. Masculinidades y salud de los hombres en la Región de las Américas. *Revista Panamericana de Salud Pública*. 2022;46: 1. <https://doi.org/10.26633/RPSP.2022.93>
 36. Del Razo-Olvera FM, Reyes-Muñoz E, Rojas-Martínez R, Guerrero-Romero F, Mehta R, Dávila-Olmedo WE, et al. Development and validation of a tool for predicting type 2 diabetes in Mexican women of reproductive age. *Endocrinol Diabetes Nutr (Engl Ed)*. 2020;67: 578–585. <https://doi.org/10.1016/j.endinu.2020.02.006>
 37. Rojas-Martínez R, Escamilla-Núñez C, Gómez-Velasco DV, Zárate-Rojas E, Aguilar-Salinas CA. Para estimar la incidencia del síndrome metabólico G colaborador de la cohorte. Diseño y validación de un score para detectar adultos con prediabetes y diabetes no diagnosticada. *Salud Publica Mex*. 2018;60: 500. <https://doi.org/10.21149/9057>
 38. Gutiérrez Zavala Á, González Avendaño MI, Grajales Castillejos O, Patiño Suárez MM, Gutiérrez Domínguez LÁ, Facultad de Ciencias Odontológicas y Salud Pública, Universidad de Ciencias y Artes de Chiapas, et al. Prevalencia de sobrepeso y obesidad en población indígena del municipio de Oxchuc, Chiapas; México. *AMU*. 2025;4: 77–83. <https://doi.org/10.31644/AMU.V04.N02.2025.A09>
 39. Sánchez-Romero LM, Sagaceta-Mejía J, Mindell JS, Passi-Solar Á, Bernabé-Ortiz A, Tolentino-Mayo L, et al. Sex differences in the secular change in waist circumference relative to body mass index in the Americas and England from 1997 to 2020.

- Epidemiology. 2023. <https://doi.org/10.1101/2023.12.10.23299756>
40. Abildgaard J, Ploug T, Al-Saoudi E, Wagner T, Thomsen C, Ewertsen C, et al. Changes in abdominal subcutaneous adipose tissue phenotype following menopause is associated with increased visceral fat mass. *Sci Rep.* 2021;11. <https://doi.org/10.1038/s41598-021-94189-2>
41. Szeliga A, Chedraui P, Meczekalski B. The Impact of the Menopausal Transition on Body Composition and Abdominal Fat Redistribution. *J Clin Med.* 2026;15. <https://doi.org/10.3390/jcm15020740>
42. Usher-Smith JA, Sharp SJ, Griffin SJ. The spectrum effect in tests for risk prediction, screening, and diagnosis. *BMJ.* 2016;353. <https://doi.org/10.1136/bmj.i3139>
43. Morales-Ortiz K, Terán-Avedaño K, Urrutia-Villanueva N, Mardones-Leiva K, Vergara-Maldonado C. Masculinidad hegemónica en la salud sexual y reproductiva: prácticas y creencias de hombres jóvenes en Chile. *MatActual.* 2021; 12. <https://doi.org/10.22370/revmat3.2021.2859>
44. Mora-Tordecillas JE de J, Ramírez-Llerena E, Benítez-Izquierdo MC, Hernández-Dáger L, Ibarra-Cervantes LDJ. Pobreza y derechos humanos de las mujeres latinoamericanas, entre 2018 y 2022: casos de México y Colombia. *Saber Cienc Lib.* 2025;19: 21–59. <https://revistas.unilibre.edu.co/index.php/saber/issue/view/749> <https://doi.org/10.18041/2382-3240/saber.2024v19n1.11355>
45. Herrera-Medina N, Espinoza-Navarro O. Aspectos Éticos a Considerar en Investigaciones con Población Indígena. *Int J Morphol.* 2025;43: 930–932. <https://doi.org/10.4067/S0717-95022025000300930>

Exactitud diagnóstica del cuestionario FINDRISC para detección de diabetes mellitus tipo 2 en población de la Sierra Nororiental del Estado de Puebla, México

RESUMEN

INTRODUCCIÓN El cuestionario *Finnish Diabetes Risk Score* es una herramienta ampliamente utilizada para el tamizaje de diabetes mellitus tipo 2; sin embargo, su evaluación en poblaciones indígenas es limitada. Este estudio evaluó la exactitud diagnóstica de este instrumento para la detección de casos compatibles con diabetes mellitus tipo 2 en población totonaca del estado de Puebla, México, y su desempeño al estratificar por sexo.

MÉTODOS Se efectuó un estudio transversal de exactitud diagnóstica con recolección prospectiva, realizado en 148 adultos (de 18 años o más) sin diagnóstico previo de diabetes mellitus tipo 2. Los participantes fueron residentes de comunidades de la Sierra Nororiental de Puebla y reclutados mediante muestreo por conveniencia con inclusión consecutiva durante una feria de salud. El cuestionario *Finnish Diabetes Risk Score* se utilizó como prueba índice, evaluado mediante dos puntos de corte (igual o superior a 12 e igual o superior a 15), y la hemoglobina glucosilada (igual o superior a 6,5%) como estándar de referencia. Ambas pruebas fueron aplicadas de forma independiente y con cegamiento entre evaluadores. Se estimaron sensibilidad, especificidad, valores predictivos, exactitud diagnóstica y área bajo la curva, mediante análisis global y estratificado por sexo.

RESULTADOS Con un valor de corte ≥ 12 , el *Finnish Diabetes Risk Score* mostró una sensibilidad general del 76,4% (intervalo de confianza: 95%; 66,6 a 84,0), especificidad de 55,9% (intervalo de confianza: 95%; 43,3 a 67,9), exactitud diagnóstica de 68,2% (intervalo de confianza: 95%; 60,4 a 75,2) y área bajo la curva de 0,64 (intervalo de confianza: 95%; 0,54 a 0,73) al punto de corte igual o superior a 12. En mujeres, la sensibilidad alcanzó 85,2% (intervalo de confianza: 95%; 73,4 a 92,3) con área bajo la curva de 0,71 (intervalo de confianza: 95%; 0,60 a 0,81), mientras que en hombres la capacidad discriminatoria fue limitada.

CONCLUSIONES El *Finnish Diabetes Risk Score* mostró capacidad discriminatoria moderada para la detección de casos compatibles con diabetes mellitus tipo 2 en población indígena totonaca, con desempeño aceptable en mujeres, utilizando el punto de corte igual o superior a 12. Estos hallazgos respaldan su uso como herramienta inicial de tamizaje en entornos con acceso limitado a pruebas confirmatorias y destacan la necesidad de validación en contextos específicos.



This work is licensed under a Creative Commons Attribution 4.0 International License.