

## Living FRIendly Summaries of the Body of Evidence using Epistemonikos (FRISBEE)

Medwave 2015; 15(Suppl 1):e6202 doi: 10.5867/medwave.2015.6202

# Should we add vancomycin antibiotic powder to post operative infection prevention in spine surgery? - First update

**Authors:** Mario López[1,3], Marcelo Molina[2,3]

**E-mail:** [mmolinas@med.puc.cl](mailto:mmolinas@med.puc.cl)

**Citation:** López M, Molina M. Should we add vancomycin antibiotic powder to post operative infection prevention in spine surgery? - First update. *Medwave* 2015; 15(Suppl 1):e6202 doi: 10.5867/medwave.2015.6202

**Publication date:** 5/8/2015

### Abstract

This Living FRISBEE (Living FRIendly Summary of the Body of Evidence using Epistemonikos) is an update of the summary published in June 2015, based on the detection of a new systematic review not identified in the previous version.

### About the update

This article updates the June 2015 Living FRISBEE (Living FRISBEE: Living FRIendly Summary of the Body of Evidence using Epistemonikos) (doi: 10.5867/medwave.2015.6160), based on the detection of a new systematic review not identified in the previous version. [1], which did not include any primary study not previously retrieved.

The new evidence incorporated in this summary does not lead to changes on the certainty of the evidence, the estimate of the magnitude of effects, the key messages or the considerations for decision-making.

### Problem

The infection rate after spinal surgery ranges from 0.5 to 12 %. For decades, efforts have been made in order to implement different measures to reduce this risk and thus improve surgical outcomes. Adding intrawound vancomycin powder could decrease the risk of infection and associated complications.

### Methods

We used Epistemonikos database, which is maintained by screening more than 30 databases, to identify systematic reviews and their included primary studies. With this information we generated a structured summary using a pre-established format, which includes key messages, a summary of the body of evidence (presented as an evidence matrix in Epistemonikos), meta-analysis of the total of studies, a summary of findings table following the GRADE approach and a table of other considerations for decision-making.

#### Key messages

- Vancomycin powder probably does not decrease surgical site infection in low-risk spinal surgery.
- There is uncertainty about the effects of vancomycin powder in populations or surgeries with a higher risk because the certainty of the evidence is very low.

### About the body of evidence for this question

What is the evidence. See evidence matrix in Epistemonikos later	We found seven systematic reviews [1],[2],[3],[4],[5],[6],[7] that consider 16 primary studies (reported in 17 references) [8],[9],[10],[11],[12],[13],[14],[15],[16],[17],[18],[19],[20],[21],[22],[23],[24]. Including only one randomized controlled trial [22].
What types of patients were included	The 16 studies included adults; three studies included posterior cervical surgery [8],[16],[20],[24], six studies (including the only randomized trial) cervical and posterior thoracolumbar surgery [9],[12],[15],[18],[21],[22], four posterior thoracolumbar surgery [13],[14],[19],[23], one posterior lumbar [17] and two studies did not specify the type of surgery [10],[11]. Three studies (including the only randomized trial) analyzed separately instrumented and non-instrumented surgery [11],[17],[22].
What types of interventions were included	The intervention was vancomycin powder. Nine studies (including the randomized) administered one gram of vancomycin powder [8],[9],[10],[15],[17],[20],[21],[22],[23], three studies used two grams [13],[14],[19], two studies 0.5 to 2 grams [11],[12], one study one to two grams [18] and one study 500 mg [16],[24]. All studies compared against standard treatment which corresponds to intravenous ceftazolin.
What types of outcomes were measured	Risk of infection, Staphylococcus aureus infection, pseudarthrosis.

### Summary of findings

The information on the effects of vancomycin powder on the surgical site is based on one randomized controlled trial including 907 patients. We conducted an evaluation of the certainty of the evidence coming from 15 non-randomized studies, which produced lower certainty than the only randomized trial. So, we considered it for the formulation of key messages and considerations for decision-making, but not for the summary of findings table.

- Vancomycin powder probably does not decrease surgical site infection in low-risk spinal surgery. The certainty of the evidence is moderate.

<b>Vancomycin powder in spinal surgery</b>				
<b>Patients</b>	spinal surgery (low-risk of infection)			
<b>Intervention</b>	vancomycin powder plus intravenous antibiotic prophylaxis			
<b>Comparison</b>	only intravenous antibiotic prophylaxis			
Outcomes	Absolute effect*		Relative effect (95% CI)	Certainty of the evidence (GRADE)
	WITH intravenous antibiotics	WITH vancomycine powder		
	Difference: patients per 1000			
Surgical site infection	17 per 1000	16 per 1000	RR 0.96 (0.35 a 2.62)	⊕⊕⊕○ <sup>1,2</sup> Moderate
	Difference: 1 less per 1000 (Margin of error: 11 less to 27 more)			
RR: Risk ratio Margin of error = 95% confidence interval (CI). GRADE: evidence grades of the GRADE Working Group (see later in this article)				
* The risk <b>WITH INTRAVENOUS ANTIBIOTICS</b> is based on the risk in the control group of the trials. The risk <b>WITH ADDED VANCOMYCIN POWDER</b> (and its margin of error) is calculated from relative effect (and its margin of error)				
<sup>1</sup> We downgraded the certainty of the evidence for risk of bias because allocation concealment and blinding were not clear.				
<sup>2</sup> Even though the study used generic vancomycin which can be considered as a reason of indirectness we did not downgrade the certainty of the evidence for this issue assuming the appropriate safeguards were taken.				

### About the certainty of the evidence (GRADE)\*

⊕⊕⊕⊕

**High:** This research provides a very good indication of the likely effect. The likelihood that the effect will be substantially different† is low.

⊕⊕⊕○

**Moderate:** This research provides a good indication of the likely effect. The likelihood that the effect will be substantially different† is moderate

⊕⊕○○

**Low:** This research provides some indication of the likely effect. However, the likelihood that it will be substantially different† is high.

⊕○○○

**Very low:** This research does not provide a reliable indication of the likely effect. The likelihood that the effect will be substantially different† is very high.

\*This concept is also called 'quality of the evidence' or 'confidence in effect estimates'.

† Substantially different = a large enough difference that it might affect a decision.

---

## Other considerations for decision-making

---

### To whom this evidence does and does not apply

---

- This evidence applies to adult patients submitted to a wide range of spinal surgeries. However, given the more reliable information comes from a single randomized trial with a relatively low risk of infection, it must be applied cautiously to patients or surgeries with a higher risk. In patients and surgeries described in the observational studies we cannot tell if vancomycin decreases risk of infection because the certainty of the evidence is very low.
  - This evidence does not apply to pediatric population which constitutes a different group of surgeries, with distinctive characteristics.
- 

### About the outcomes included in this summary

---

- The selected outcome is surgical site infection, which constitutes the critical outcome for decision-making according to the main guideline and the judgment of the authors of this summary.
- 

### Balance between benefits and risks, and certainty of the evidence

---

- Considering there is no benefit with moderate certainty it is unlikely that the benefit/risk ratio for this intervention is favourable.
  - Given the higher risk of infection in some cases, and the relatively low-cost and safety of the intervention, some clinicians might be inclined to use it despite existing uncertainty in immunosuppressed patients, long-term steroid use, associated neuromuscular disease, previous infected surgery, long instrumented arthrodeses, double approach (anterior and posterior), revision surgery and prolonged surgery.
- 

### Resource considerations

---

- Considering the probable absence of benefit, it is unlikely it constitutes a cost-effective intervention, but it is not possible to make any inference for patients or surgeries with a higher risk of infection based on the analysed studies.
- 

### Differences between this summary and other sources

---

- Our summary is in agreement with the more recent systematic review that warns the evidence is weak and more information is needed before adopting this intervention [3]. The other reviews identified provide a more favourable conclusion because they did not include the randomized study or did not take into consideration the differences in study designs when making conclusions.
  - Our summary is in partial agreement with the main guideline in this area (North American Spine Society) [25] which suggests using vancomycin powder in prolonged surgeries, instrumented surgeries or comorbidities such as diabetes, neuromuscular disease and neurological damage. This guideline does not include the randomized trial, since it incorporated evidence up to June 2011.
- 


### Could this evidence change in the future?

---

- The probability of future evidence changing what we know is low, because of the level of certainty of the evidence.
  - Future controlled studies in patients or surgeries with a higher risk of infection have a high likelihood of changing what we know.
-

## How we conducted this summary

Using automated and collaborative means, we compiled all the relevant evidence for the question of interest and we present it as a matrix of evidence.



Starting from any systematic review, Epistemonikos builds a matrix based on existing connections in the database.

The author of the matrix can select relevant information for a specific health question (typically in PICO format) in order to display the information set for the question.

The rows represent systematic reviews that share at least one primary study, and columns display the studies.

The boxes in green correspond to studies included in the respective reviews.

Follow the link to access the **interactive version**: [Vancomycin powder vs endovenous antibiotic prophylaxis to avoid surgical site infection in patients with spine surgery](#)

## Notes

The upper portion of the matrix of evidence will display a warning of “new evidence” if new systematic reviews are published after the publication of this summary. Even though the project considers the periodical update of these summaries, users are invited to comment in *Medwave* or to contact the authors through email if they find new evidence and the summary should be updated earlier. After creating an account in Epistemonikos, users will be able to save the matrixes and to receive automated notifications any time new evidence potentially relevant for the question appears. The details about the methods used to produce these summaries are described here <http://dx.doi.org/10.5867/medwave.2014.06.5997>.

Epistemonikos foundation is a non-for-profit organization aiming to bring information closer to health decision-makers with technology. Its main development is Epistemonikos database ([www.epistemonikos.org](http://www.epistemonikos.org)).

These summaries follow a rigorous process of internal peer review.

## Conflicts of interest

The authors do not have relevant interests to declare.

## References

- Xiong L, Pan Q, Jin G, Xu Y, Hirche C. Topical intrawound application of vancomycin powder in addition to intravenous administration of antibiotics: A meta-analysis on the deep infection after spinal surgeries. *Orthop Traumatol Surg Res.* 2014 Nov;100(7):785-9. | [CrossRef](#) | [PubMed](#) |
- Khan NR, Thompson CJ, DeCuypere M, Angotti JM, Kalobwe E, Muhlbauer MS, et al. A meta-analysis of spinal surgical site infection and vancomycin powder. *J Neurosurg Spine.* 2014 Dec;21(6):974-83. | [CrossRef](#) | [PubMed](#) |
- Kang DG, Holekamp TF, Wagner SC, Lehman RA Jr. Intracavitary vancomycin powder for the prevention of surgical site infection in spine surgery: a systematic



- literature review. *Spine J.* 2015 Apr 1;15(4):762-70. | [CrossRef](#) | [PubMed](#) |
4. Alcalá-Cerra G, Paternina-Cacedo AJ, Moscote-Salazar LR, Gutiérrez-Paternina JJ, Niño-Hernández LM. [Application of vancomycin powder into the wound during spine surgery: systematic review and meta-analysis]. *Rev Esp Cir Ortop Traumatol.* 2014 May-Jun;58(3):182-91. | [CrossRef](#) | [PubMed](#) |
  5. Chiang HY, Herwaldt LA, Blevins AE, Cho E, Schweizer ML. Effectiveness of local vancomycin powder to decrease surgical site infections: a meta-analysis. *Spine J.* 2014 Mar 1;14(3):397-407. | [CrossRef](#) | [PubMed](#) |
  6. Bakhsheshian J, Dahdaleh NS, Lam SK, Savage JW, Smith ZA. The use of vancomycin powder in modern spine surgery: systematic review and meta-analysis of the clinical evidence. *World Neurosurg.* 2015 May;83(5):816-23. | [CrossRef](#) | [PubMed](#) |
  7. Kanj WW, Flynn JM, Spiegel DA, Dormans JP, Baldwin KD. Vancomycin prophylaxis of surgical site infection in clean orthopedic surgery. *Orthopedics.* 2013 Feb;36(2):138-46. | [CrossRef](#) | [PubMed](#) |
  8. Caroom C, Tullar JM, Benton EG Jr, Jones JR, Chaput CD. Intrawound vancomycin powder reduces surgical site infections in posterior cervical fusion. *Spine (Phila Pa 1976).* 2013 Jun 15;38(14):1183-7. | [CrossRef](#) | [PubMed](#) |
  9. Godil SS, Parker SL, O'Neill KR, Devin CJ, McGirt MJ. Comparative effectiveness and cost-benefit analysis of local application of vancomycin powder in posterior spinal fusion for spine trauma: clinical article. *J Neurosurg Spine.* 2013 Sep;19(3):331-5. | [CrossRef](#) | [PubMed](#) |
  10. Rahman RKK, Lenke LG, Bridwell KH, Buchowski J, Dickson DD, Aleem A, et al. Intrawound vancomycin powder lowers the acute deep wound infection rate in adult spinal deformity patients. *Spine: Affiliated Society Meeting Abstracts.* 2011;(suppl 2011):73. | [Link](#) |
  11. Lai S-M, Heller A. The effect of intrawound vancomycin powder on surgical site infections in posterior instrumented spinal arthrodeses [Submitted to the graduate degree program in Clinical Research in partial fulfillment of the requirements for the degree of Master of Science ] University of Kansas, 2012. | [Link](#) |
  12. Heller A, McIff TE, Lai SM, Burton DC. Intrawound Vancomycin Powder Decreases Staphylococcal Surgical Site Infections Following Posterior Instrumented Spinal Arthrodesis. *J Spinal Disord Tech.* 2013 Oct 30. [Epub ahead of print] . | [PubMed](#) |
  13. Sweet FA, Roh M, Sliva C. Intrawound application of vancomycin for prophylaxis in instrumented thoracolumbar fusions: efficacy, drug levels, and patient outcomes. *Spine (Phila Pa 1976).* 2011 Nov 15;36(24):2084-8. | [CrossRef](#) | [PubMed](#) |
  14. Theologis AA, Demirkiran G, Callahan M, Pekmezci M, Ames C, Deviren V. Local intrawound vancomycin powder decreases the risk of surgical site infections in complex adult deformity reconstruction: a cost analysis. *Spine (Phila Pa 1976).* 2014 Oct 15;39(22):1875-80. | [CrossRef](#) | [PubMed](#) |
  15. Kim HS, Lee SG, Kim WK, Park CW, Son S. Prophylactic intrawound application of vancomycin powder in instrumented spinal fusion surgery. *Korean J Spine.* 2013 Sep;10(3):121-5. | [CrossRef](#) | [PubMed](#) |
  16. Pahys JM, Pahys JR, Cho SK, Kang MM, Zebala LP, Hawasli AH, et al. Methods to decrease postoperative infections following posterior cervical spine surgery. *J Bone Joint Surg Am.* 2013 Mar 20;95(6):549-54. | [CrossRef](#) | [PubMed](#) |
  17. Strom RG, Pacione D, Kalthorn SP, Frempong-Boadu AK. Lumbar laminectomy and fusion with routine local application of vancomycin powder: decreased infection rate in instrumented and non-instrumented cases. *Clin Neurol Neurosurg.* 2013 Sep;115(9):1766-9. | [CrossRef](#) | [PubMed](#) |
  18. Hill BW, Emohare O, Song B, Davis R, Kang MM. The use of vancomycin powder reduces surgical reoperation in posterior instrumented and noninstrumented spinal surgery. *Acta Neurochir (Wien).* 2014 Apr;156(4):749-54. | [CrossRef](#) | [PubMed](#) |
  19. Martin JR, Adogwa O, Brown CR, Bagley CA, Richardson WJ, Lad SP, et al. Experience with intrawound vancomycin powder for spinal deformity surgery. *Spine (Phila Pa 1976).* 2014 Jan 15;39(2):177-84. | [CrossRef](#) | [PubMed](#) |
  20. Strom RG, Pacione D, Kalthorn SP, Frempong-Boadu AK. Decreased risk of wound infection after posterior cervical fusion with routine local application of vancomycin powder. *Spine (Phila Pa 1976).* 2013 May 20;38(12):991-4. | [CrossRef](#) | [PubMed](#) |
  21. O'Neill KR, Smith JG, Abtahi AM, Archer KR, Spengler DM, McGirt MJ, et al. Reduced surgical site infections in patients undergoing posterior spinal stabilization of traumatic injuries using vancomycin powder. *Spine J.* 2011 Jul;11(7):641-6. | [CrossRef](#) | [PubMed](#) |
  22. Tubaki VR, Rajasekaran S, Shetty AP. Effects of using intravenous antibiotic only versus local intrawound vancomycin antibiotic powder application in addition to intravenous antibiotics on postoperative infection in spine surgery in 907 patients. *Spine (Phila Pa 1976).* 2013 Dec 1;38(25):2149-55. | [CrossRef](#) | [PubMed](#) |
  23. Emohare O, Ledonio CG, Hill BW, Davis RA, Polly DW Jr, Kang MM. Cost savings analysis of intrawound vancomycin powder in posterior spinal surgery. *Spine J.* 2014 Nov 1;14(11):2710-5. | [CrossRef](#) | [PubMed](#) |
  24. Pahys JM, Pahys JR, Cho SK, Kang MM, Zebala LP, Hawasli AH, et al. Methods to decrease postoperative infections following posterior cervical spine surgery. *J Bone Joint Surg Am.* 2013 Mar 20;95(6):549-54. | [CrossRef](#) | [PubMed](#) |
  25. Shaffer WO, Baisden JL, Fernand R, Matz PG; North American Spine Society. An evidence-based clinical guideline for antibiotic prophylaxis in spine surgery. *Spine J.* 2013 Oct;13(10):1387-92. | [CrossRef](#) | [PubMed](#) |

**Author address:**

**[1]** Facultad de Medicina  
Pontificia Universidad Católica de Chile  
Lira 63  
Santiago Centro  
Chile



Esta obra de Medwave está bajo una licencia Creative Commons Atribución-No Comercial 3.0 Unported. Esta licencia permite el uso, distribución y reproducción del artículo en cualquier medio, siempre y cuando se otorgue el crédito correspondiente al autor del artículo y al medio en que se publica, en este caso, Medwave.